

Key Findings

- The number of samples tested (176) has continued to increase this month. The increase is due to the addition of a new and growing service with the Harm Reduction Team (HaRT) at UBCO.
- The percentage of opioids testing positive for benzodiazepines in the Interior Health region (36%) has remained consistent from last month but has been on upward trend since September (6%).
- There were 8 drug checking alerts in Interior Health for this month highlighting drugs adulterated with benzodiazepines.







Number of samples tested with fentanyl present

101 Opioid samples

95 FENTANYL POSITIVE



6 FENTANYL NEGATIVE

36 Stimulant samples

34 FENTANYL NEGATIVE



2 FENTANYL POSITIVE

1 Depressant samples



1 FENTANYL NEGATIVE

23 Psychedelic samples



23 FENTANYL NEGATIVE

14 Unknown samples

11 FENTANYL NEGATIVE (79%)



2 FENTANYL POSITIVE (14%)

2 Polysubstance sample




2 FENTANYL POSITIVE (100%)



PUBLIC HEALTH NOTIFICATIONS

Date & Location	Expected Drug	Drugs Detected	Fentanyl Strip	Benzo Strip	Area Purchased	Alert Message
17 February 2021- ASK Wellness, Kamloops	Percocet	Acetaminophen and fentanyl	Positive	N/A	Kamloops	Unknown consumption of fentanyl poses a high risk for opioid overdose! Check your drugs at any local SCS/OPS!
6 February 2021- UBCO Kelowna/West Kelowna	Fentanyl	Contains benzodiazepines and fentanyl	Positive	Positive	Kelowna	Check your drugs for benzodiazepines at any local SCS/OPS.
5 Februrary 2021- ANKORS, Cranbrook	Down	Contains benzodiazepines and high concentrations of fentanyl	Positive	Positive	Cranbrook	High risk of prolonged sedation and overdose- Check your drugs for benzodiazepines at any local SCS/OPS.
5 Februrary 2021- ANKORS, Cranbrook	Down	High fentanyl concentration and benzodiazepines	Positive	Positive	Cranbrook	May not respond to naloxone- continue to give breaths and seek medical attention.
5 Februrary 2021- ASK Wellness, Kamloops	Cocaine	Methamphetamine, cocaine, levamisole	N/A	N/A	Kamloops	Meth is a powerful stimulant that has longer lasting effects than cocaine. Check your drugs for unexpected substances!
4 Februrary 2021- UBCO Kelowna/West Kelwona	Fentanyl	Benzodiazepines and fentanyl	Positive	Positive	Kelowna	High risk of overdose with prolonged sedation-may not respond to Naloxone. Use at an OPS!
3 Februrary 2021- ANKORS, Castlegar	Fentanyl	Fentanyl and benzodiazepines	Positive	Positive	Castlegar	Check for benzos in your drugs at any SCS or OPS!
2 Februrary 2021- IHA Pentiction	Down	High concentration of fentanyl and benzodiazepines	Postitive	Positive	Penticton	High risk of overdose with prolonged sedationn

For information about public health alerts in the Interior Health region, please visit:



During the month of February, 36% of expected opioid samples tested positive for benzodiazepines using test strips in our partner sites in the Interior Health region (36 samples of 100 tested).

Opioid samples are tested for benzodiazepine-positivity using BTNX test strips. The results presented above are presumptive and tentative until confirmed by a laboratory.

Number of drugs checks at each site

Organization	City/Town	# of Drugs Checked
ANKORS	Cranbrook	10
ANNORS	Nelson	36
	Kamloops	45
ASK Wellness	Merritt	9
ASK Welliess	Vernon	8
	Mobile Bus	4
	Kelowna	24
UBC Okanagan	Penticton	30
	Vernon	10
Total		176

Number of samples that matched expectation

using FTIR/test strip drug checking

EXPECTED DRUG: Depressant

1 Sample Tested

EXPECTED DRUG:

Psychedelic

23 Samples Tested




Matched: 23

EXPECTED DRUG:

Stimulant

36 Samples Tested



Matched: 34 Did not match: 2 EXPECTED DRUG:

Polysubstance

2 Samples Tested



Matched: 2

EXPECTED DRUG:

Opioid

Matched: 1

101 Samples Tested



Matched: 95
Did not match: 4
Match not determined: 2

EXPECTED DRUG:

Unknown

14 Samples Tested



Match not determined: 14

Total

177 Samples Tested



Matched: 155
Did not match: 6

Match not determined: 16

Please note that the presence of the expected substance does not imply purity, as samples frequently contain adulterating cutting agents

Number of opioid samples that matched expectation

using FTIR/test strip drug checking

EXPECTED DRUG:

Fentanyl

81 Samples Tested



Matched: 81 Matched Did not

EXPECTED DRUG:

Heroin

4 Samples Tested



Matched: 3 Did not match: 1 **EXPECTED DRUG:**

Down

10 Samples Tested



Matched: 9
Did not match: 1

EXPECTED DRUG:

Pharmaceutical

6 Samples Tested



Matched: 2

Did not match: 2
Match not determined: 2

Total

101 Samples Tested



Matched: 95

Did not match: 4

Match not determined: 2

Please note that the presence of the expected substance does not imply purity, as samples frequently contain adulterating cutting agents. 'Down' can refer to any opioid drug present in any amount.

Data represented here are collected from our partner sites across the province. Drug samples are tested using the Fourier Transform Infrared (FTIR) spectrometer in combination with fentanyl test strips and benzodiazapine test strips.

There is 5% fentanyl detection limit on the FTIR spectrometer (McCrae, 2019), and a drug check on any given sample consists of <u>both</u> the FTIR <u>and</u> BTNX fentanyl immunoassay test strip testing done in combination. When applicable, BTNX benzodiazepine immunoassay test strips are also used.

BCCSU gratefully acknowledges the contributions of the following partners:









